

**Ogsiveo**<sup>™</sup>  
(nirogacestat)  
50 mg tablets

Print double-sided and fold here

# DOCTOR DISCUSSION GUIDE

A desmoid tumor diagnosis can feel overwhelming, so it is important to communicate with your doctor. This guide provides questions and helpful context to discuss with your doctor to see if OGSIVEO is right for you. You can also find information about what you might expect during treatment with OGSIVEO.

**OGSIVEO is the first and only FDA-approved treatment for adults with progressing desmoid tumors who require a medicine by mouth or injection (systemic therapy). It is not known if OGSIVEO is safe and effective in children.**

**OGSIVEO can cause serious side effects, including diarrhea, ovarian problems, liver problems, new non-melanoma skin cancers, and electrolyte (salt) problems.**

Please see Important Safety Information throughout and click here to view [full Prescribing Information including Patient Information](#), or visit [OGSIVEO.com](#).



## Who should be involved in my desmoid tumor treatment?

Consider seeking advice from a multidisciplinary care team experienced in treating desmoid tumors. They may offer expert advice about your care with OGSIVEO. This team may include a medical oncologist and sarcoma specialist, surgeon or surgical oncologist, radiologist, radiation oncologist, pathologist, nurse, nurse practitioner, and physician assistant.<sup>1,2</sup>



## What is OGSIVEO?

OGSIVEO is a prescription medication used to treat adults with progressing desmoid tumors who require a medicine by mouth or injection (systemic therapy). It is not known if OGSIVEO is safe and effective in children. Discuss your medical history with your doctor—and together, you can decide if OGSIVEO is an option for you.



## What should I consider before deciding to take OGSIVEO?

OGSIVEO is not for everyone. Tell your doctor about all of your medical conditions, including if you have liver problems, are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed. Tell your doctor about all the medicines you take. This will help you and your doctor to see if OGSIVEO may be right for you.<sup>3</sup>



## Should I be concerned about side effects with OGSIVEO?

Always talk with your doctor about possible side effects of any medication you take. It's important to know what you may experience and how your doctor may help. Learn about possible side effects of OGSIVEO in the **Important Safety Information** throughout this guide.



## How should I take OGSIVEO?

**Take OGSIVEO exactly as your healthcare provider tells you to take it.** OGSIVEO is an oral medication that comes in 50-mg tablets. The recommended dosage is 150 mg (which is three 50-mg tablets), taken 2 times a day—a **total of 6 tablets**. OGSIVEO can be taken with or without food. Swallow OGSIVEO tablets whole. Do not break, crush, or chew tablets. If you take an antacid medicine, take OGSIVEO 2 hours before or 2 hours after taking the antacid. If you vomit after taking a dose or miss a dose of OGSIVEO, take your next dose at your regular time. Do not take 2 doses of OGSIVEO to make up the dose.<sup>3</sup>



## What are the next steps with OGSIVEO?

If your doctor determines that OGSIVEO is an option for you, you can learn about support available for people taking OGSIVEO at [www.ogsiveo.com/patient-support-program](http://www.ogsiveo.com/patient-support-program).



**Tip: Print out this guide to refer to when you meet with your doctor.**

## What is OGSIVEO?

OGSIVEO is a prescription medication used to treat adults with progressing desmoid tumors who require a medicine by mouth or injection (systemic therapy). It is not known if OGSIVEO is safe and effective in children.

## Important Safety Information

**Before taking OGSIVEO tell your healthcare provider about all of your medical conditions, including if you:**

- Have liver problems.
- Are pregnant or plan to become pregnant. OGSIVEO can harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment.

### Females who are able to become pregnant:

- Your healthcare provider will give you a pregnancy test before you start treatment with OGSIVEO.
- You should use effective birth control during treatment and for 1 week after the last dose. Talk to your healthcare provider about methods that may be right for you.
- Stop taking OGSIVEO and tell your healthcare provider right away if you become pregnant.

**Males** with female partners who are able to become pregnant should use effective birth control during treatment with OGSIVEO and for 1 week after the last dose.

- Are breastfeeding or plan to breastfeed. It is not known if OGSIVEO passes into your breast milk. Do not breastfeed during treatment with OGSIVEO and for 1 week after the last dose.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You should avoid taking proton pump inhibitors (PPIs) and H2 blockers during treatment with OGSIVEO. Ask your healthcare provider if you are not sure if you take one of these medicines.

## How should I take OGSIVEO?

- Take OGSIVEO exactly as your healthcare provider tells you to take it.
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with OGSIVEO if you develop side effects.
- Take OGSIVEO 2 times a day with or without food.
- Swallow OGSIVEO tablets whole. Do not break, crush, or chew.
- If you take an antacid medicine, take OGSIVEO 2 hours before or 2 hours after the antacid.
- If you vomit after taking a dose or miss a dose of OGSIVEO, take your next dose at your regular time. Do not take 2 doses of OGSIVEO to make up the dose.

## What should I avoid while taking OGSIVEO?

Avoid eating or drinking grapefruit products, Seville oranges, and starfruit during treatment with OGSIVEO.

## What are the possible side effects of OGSIVEO?

OGSIVEO can cause serious side effects, including:

- **Diarrhea.** Diarrhea is common with OGSIVEO and may sometimes be severe. Your healthcare provider may tell you to drink more fluids or to take antidiarrheal medicines. Tell your healthcare provider right away if you have diarrhea that lasts longer than a few days and does not get better after taking antidiarrheal medicines.
- **Ovarian problems.** Females who are able to become pregnant may have ovarian problems and changes in their menstrual cycle during treatment. OGSIVEO may affect fertility which may affect your ability to have a child. Tell your healthcare provider if you have any changes in your menstrual cycle or hot flashes, night sweats, or vaginal dryness during treatment.

Print double-sided and fold here



## Preparing for your appointment:

- Consider keeping track of your symptoms and letting your doctor or care team know if anything has changed
- Bring a list of any medications or supplements you take and their doses
- Consider bringing a family member or friend with you for support during the visit
- Bring your insurance card, ID, and medical records if your doctor doesn't already have them

- **Liver problems.** OGSIVEO can increase liver enzymes. Your healthcare provider will do blood tests to check your liver function before you start and during treatment with OGSIVEO.
- **New non-melanoma skin cancers.** Your healthcare provider will do skin exams before and during treatment with OGSIVEO if you are at risk for skin cancer. Tell your healthcare provider if you have any new or changing skin lesions.
- **Electrolyte (salt) problems.** Your healthcare provider will do blood tests to check your phosphate and potassium levels during treatment and may give you medicines to treat low phosphate or low potassium if needed. Tell your healthcare provider if you develop any muscle pain or weakness.

Tell your healthcare provider right away if you have any side effect that bothers you or that does not go away. Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with OGSIVEO.

## The most common side effects of OGSIVEO are:

- rash
- headache
- hair loss
- nausea
- stomach (abdominal) pain
- upper respiratory infection
- tiredness
- mouth sores
- cough
- shortness of breath

OGSIVEO can affect fertility in females and males, which may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of OGSIVEO.

Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-FDA-1088.

**Please see Important Safety Information throughout and click here to view full Prescribing Information including Patient Information, or visit [OGSIVEO.com](https://ogsiveo.com).**

**References:** 1. Singhi E, Feldman J, West H. The multidisciplinary cancer conference. *JAMA Oncology*. Published online: December 22, 2022. doi:10.1001/jamaoncol.2022.4924. 2. Cancer.Net. The oncology team. Accessed October 16, 2023. <https://www.cancer.net/navigating-cancer-care/cancer-basics/cancer-care-team/oncology-team> 3. OGSIVEO. Prescribing information. SpringWorks Therapeutics, Inc.

**The information and questions provided in this Guide serve as suggestions only to prompt discussions that you may wish to have with your care team, but they are not intended to be medical advice. Always follow the medical advice of your care team regarding any medical condition or treatment.**

