



Actor portrayal. Individual results may vary.

TREATMENT TRACKER

A resource to help you track your journey with OGSIVEO

The **OGSIVEO Treatment Tracker** is a tool for adults with progressing desmoid tumors who have been prescribed OGSIVEO. This simple, easy-to-use document can be used to track symptoms, possible side effects, medications, supplements, and recommendations from your care team—all in one place. Logging these details can help you share your experience during treatment with your care team.

Tracking Symptoms and Side Effects

Tracking the frequency and severity of your symptoms and possible side effects can help your care team support you during treatment.

What is OGSIVEO?

OGSIVEO is a prescription medication used to treat adults with progressing desmoid tumors who require a medicine by mouth or injection (systemic therapy). It is not known if OGSIVEO is safe and effective in children.¹

Important Safety Information

Before taking OGSIVEO tell your healthcare provider about all of your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant. OGSIVEO can harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with OGSIVEO

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.

Ogsiveo[®]
(nirogacestat)
150 mg & 100 mg tablets

Overview of the OGSIVEO Treatment Tracker

Care Team Information..... page 3

You can fill out this page with the names and contact information of your care team, which may include your medical oncologist, nurses, and any other healthcare professional assisting you during your treatment.

Appointments and Treatment Goals..... page 4

This page can help you keep track of appointments and capture any goals, notes, or important takeaways from your visits.

Current Medications, Vitamins, and Supplements..... page 5

You can use this page to list all the medicines you currently take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Symptoms and Side Effects Trackers..... pages 6-7

You can use the charts provided to track any desmoid tumor symptoms or potential side effects you may experience during treatment with OGSIVEO. This information can be helpful to share with your care team.

Consider documenting details on the days you experience symptoms and possible side effects.

Medications and Other Approaches for Managing Side Effects..... page 8

You can use this page to list any medications or other treatment approaches recommended by your care team to help you manage side effects you might experience with OGSIVEO.

This document can be saved or reprinted to use on a recurring basis.



Need assistance getting started on or staying on track with OGSIVEO?

SpringWorks CareConnections™ is a free, personalized patient support program to help you navigate your insurance coverage and answer questions about your treatment.

For questions and further information on the support available through SpringWorks CareConnections, please call 844-CARES-55 (844-227-3755), Monday – Friday 8 AM – 10 PM ET or visit www.springworkstxcare.com

Please see Important Safety Information and full Prescribing Information and Patient Information for OGSIVEO.



Care Team Information

You can fill out this page with the names and contact information of your care team, which may include your medical oncologist, nurses, and any other healthcare professional assisting you during your treatment.

Medical Oncologist

Name:

Email:

Phone:

Other Specialty

Name:

Email:

Phone:

Nurse

Name:

Email:

Phone:

Nurse

Name:

Email:

Phone:

Other

Name:

Email:

Phone:

Other

Name:

Email:

Phone:

**Tell your care team about any possible side effects you experience while taking OGSIVEO.
You may report side effects to the FDA at 1-800-FDA-1088.**

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.



Tracking Appointments and Treatment Goals

Month/Year:

This page can help you keep track of appointments and capture any goals, notes, or important takeaways from your visits.

Date:

Goals:

Who I met with:

Notes:

Date:

Goals:

Who I met with:

Notes:

Date:

Goals:

Who I met with:

Notes:

Date:

Goals:

Who I met with:

Notes:

If you think your symptoms might be changing, it is a good idea to reach out to your care team so you can let them know what you're experiencing.

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.



Current Medications, Vitamins, and Supplements

It is important to tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. You can list them in the space below. Consider including what the medication, vitamin, or supplement is intended for, when you started taking it, and the dosage you take.

- You should avoid taking proton pump inhibitors (PPIs) and H2 blockers during treatment with OGSIVEO
- Ask your doctor if you are not sure if you take one of these medicines
- Avoid eating or drinking grapefruit products, Seville oranges, and starfruit during treatment with OGSIVEO

Medicines, Vitamins, and Supplements:

Notes:

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.



Symptom Tracker

Desmoid tumors can affect each person differently depending on where the tumor is in the body, how fast it grows, and other factors. Some common desmoid tumor symptoms include limited movement, deformity, pain, or muscle problems.¹ These are not all the possible symptoms of desmoid tumors, and OGSIVEO may not help to treat all or any of these symptoms. For a more comprehensive list of symptoms, please consult your care team.

Symptoms	Week beginning:	Week beginning:	Week beginning:	Week beginning:
<p>Fill in any desmoid tumor symptoms you may be experiencing.</p> <p>Consider noting any changes in the severity and frequency of your symptoms during treatment with OGSIVEO.</p> <p>For example, it may be helpful to rate your level of pain from 0 (no pain) to 10 (pain as bad as you can imagine).</p> <p>This information can be helpful to share with your care team.</p>				

These pages can be saved or reprinted to use on a recurring basis.

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.



Side Effect Tracker

Dosage:

The most common side effects of OGSIVEO are rash, nausea, tiredness, mouth sores, headache, stomach (abdominal) pain, cough, hair loss, upper respiratory infection, and shortness of breath.

Ovarian problems: Females who are able to become pregnant may have ovarian problems and changes in their menstrual cycle during treatment with OGSIVEO. OGSIVEO may affect fertility which may affect your ability to have a child. Tell your healthcare provider if you have any changes in your menstrual cycle or any hot flashes, night sweats, or vaginal dryness during treatment with OGSIVEO.

These are not all of the possible side effects of OGSIVEO. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Possible Side Effects	Week beginning:	Week beginning:	Week beginning:	Week beginning:
<p>Fill in any possible side effects you may be experiencing.</p> <p>It may be helpful to include details about how possible side effects impact your body and your day-to-day life. Consider recording the following details about possible side effects:</p> <ul style="list-style-type: none">– What does it look and feel like?– Which part of your body is affected?– When did it occur?– Is it ongoing?– How often do you experience it (eg, constantly, daily, weekly)?– Has it resolved? If so, when? <p>This information can be helpful to share with your care team.</p>				

These pages can be saved or reprinted to use on a recurring basis.

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.

Ogsiveo[®]
(nirogacestat)
150 mg & 100 mg tablets

Your doctor may recommend medications or other approaches to help you manage possible side effects

You can use this page to list any medications or other treatment approaches recommended by your care team to help you manage side effects you might experience with OGSIVEO. If listing a medication, supplement, and/or other treatment approach, consider including what it is intended for, when you started taking it, and the dosage you take (if applicable).

Medications and/or Supplements:

Other approaches:

Notes:

Please see [Important Safety Information](#) and full [Prescribing Information and Patient Information](#) for OGSIVEO.



Important Safety Information for OGSIVEO

What is OGSIVEO?

OGSIVEO is a prescription medication used to treat adults with progressing desmoid tumors who require a medicine by mouth or injection (systemic therapy). It is not known if OGSIVEO is safe and effective in children.

Important Safety Information

Before taking OGSIVEO tell your healthcare provider about all of your medical conditions, including if you:

- Have liver problems.
- Are pregnant or plan to become pregnant. OGSIVEO can harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment.

Females who are able to become pregnant:

- Your healthcare provider will give you a pregnancy test before you start treatment with OGSIVEO.
- You should use effective birth control during treatment and for 1 week after the last dose. Talk to your healthcare provider about methods that may be right for you.
- Stop taking OGSIVEO and tell your healthcare provider right away if you become pregnant.

Males with female partners who are able to become pregnant should use effective birth control during treatment with OGSIVEO and for 1 week after the last dose.

- Are breastfeeding or plan to breastfeed. It is not known if OGSIVEO passes into your breast milk. Do not breastfeed during treatment with OGSIVEO and for 1 week after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You should avoid taking proton pump inhibitors (PPIs) and H2 blockers during treatment with OGSIVEO. Ask your healthcare provider if you are not sure if you take one of these medicines.

How should I take OGSIVEO?

- Take OGSIVEO exactly as your healthcare provider tells you to take it.
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with OGSIVEO if you develop side effects.
- Take OGSIVEO 2 times a day with or without food.
- Swallow OGSIVEO tablets whole. Do not break, crush, or chew.
- If you take an antacid medicine, take OGSIVEO 2 hours before or 2 hours after the antacid.
- If you vomit after taking a dose or miss a dose of OGSIVEO, take your next dose at your regular time. Do not take 2 doses of OGSIVEO to make up the dose.

What should I avoid while taking OGSIVEO?

Avoid eating or drinking grapefruit products, Seville oranges, and starfruit during treatment with OGSIVEO.

Please see full [Prescribing Information and Patient Information](#) for OGSIVEO.



Important Safety Information for OGSIVEO (continued)

What are the possible side effects of OGSIVEO?

OGSIVEO can cause serious side effects, including:

- **Diarrhea.** Diarrhea is common with OGSIVEO and may sometimes be severe. Your healthcare provider may tell you to drink more fluids or to take antidiarrheal medicines. Tell your healthcare provider right away if you have diarrhea that lasts longer than a few days and does not get better after taking antidiarrheal medicines.
- **Ovarian problems.** Females who are able to become pregnant may have ovarian problems and changes in their menstrual cycle during treatment. OGSIVEO may affect fertility which may affect your ability to have a child. Tell your healthcare provider if you have any changes in your menstrual cycle or hot flashes, night sweats, or vaginal dryness during treatment.
- **Liver problems.** OGSIVEO can increase liver enzymes. Your healthcare provider will do blood tests to check your liver function before you start and during treatment with OGSIVEO.
- **New non-melanoma skin cancers.** Your healthcare provider will do skin exams before and during treatment with OGSIVEO if you are at risk for skin cancer. Tell your healthcare provider if you have any new or changing skin lesions.
- **Electrolyte (salt) problems.** Your healthcare provider will do blood tests to check your phosphate and potassium levels during treatment and may give you medicines to treat low phosphate or low potassium if needed. Tell your healthcare provider if you develop any muscle pain or weakness.

Tell your healthcare provider right away if you have any side effect that bothers you or that does not go away. Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with OGSIVEO.

The most common side effects of OGSIVEO are:

- rash
- stomach (abdominal) pain
- nausea
- cough
- tiredness
- hair loss
- mouth sores
- upper respiratory infection
- headache
- shortness of breath

OGSIVEO can affect fertility in females and males, which may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of OGSIVEO. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-FDA-1088.

Please see full [Prescribing Information and Patient Information](#) for OGSIVEO.

Reference: 1. Gounder MM, Maddux L, Paty J, Atkinson TM. Prospective development of a patient-reported outcomes instrument for desmoid tumors or aggressive fibromatosis. *Cancer*. 2020;126(3):531-539.



© 2024 SpringWorks Therapeutics, Inc. All rights reserved. C_OGS_US_0263 05/24
OGSIVEO is a registered trademark and SpringWorks CareConnections is a trademark of SpringWorks Therapeutics, Inc.

